UNCLASSIFIED

AD NUMBER

AD819938

LIMITATION CHANGES

TO:

Approved for public release; distribution is unlimited. Document partially illegible.

FROM:

Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; SEP 1967. Other requests shall be referred to Air Force Technical Application Center, Washington, DC 20330. Document partially illegible. This document contains export-controlled technical data.

AUTHORITY

usaf ltr, 28 feb 1972

SPATIAL CORRELATION OF AMPLITUDE ANOMALIES

7 September 1967

Prepared For

AIR FORCE TECHNICAL APPLICATIONS CENTER Washington, D. C.

By

F. A. Klappenberger TELEDYNE, INC.

Under

Project VELA UNIFORM

Sponsored By

ADVANCED RESEARCH PROJECTS AGENCY Nuclear Test Detection Office ARPA Order No. 624



BEST AVAILABLE COPY

SPATIAL CORRELATION OF AMPLITUDE ANOMALIES SEISMIC DATA LABORATORY REPORT NO. 195

AFTAC Project No.: VELA T/6702

Project Title: Seismic Data Laboratory

ARPA Order No.: 624

ARPA Program Code No.: 5810

Name of Contractor: TELEDYNE, INC.

Contract No.: F 33657-67-C-1313

Date of Contract: 2 March 1967

Amount of Contract: \$ 1, 736,617

Contract Expiration Date: 1 March 1968

Project Manager: William C. Dean (703) 836-7644

P. O. Box 334, Alexandria, Virginia

AVAILABILITY

This document is subject to special export controls and each transmittal to foreign governments or foreign national may be made only with prior approval of Chief, AFTAC.

This research was supported by the Advanced Research Projects Agency, Nuclear Test Detection Office, under Project VELA-UNIFORM and accomplished under the technical direction of the Air Force Technical Applications Center under Contract F 33657-67-C-1313.

Neither the Advanced Research Projects Agency nor the Air Force Technical Applications Center will be responsibile for information contained herein which may have been supplied by other organizations or contractors, and this document is subject to later revision as may be necessary.

TABLE OF CONTENTS

			Page	No
	ABS	TRACT		
1.	INT	RODUCTION	1	
2.	SPA	TIAL CORRELATIONS OVER LASA	2	
	Α.	LASA Spatial Correlations Using Average Log of Normalized Peak-To-Peak Amplitudes	3	
	В.	LASA Spatial Correlations of Single Events	4	
3.	SPA	TIAL CORRELATIONS OVER LASA SUBARRAYS	4	
	Α.	Subarray Spatial Correlations for Single Events	5	
	В.	Testing For Correlations Among Three Colombia Earthquakes	6	
	c.	Subarray Spatial Correlations For Two Shots	6	
	D.	Testing For Correlation Between Two Kazakh Shots	7	
4.	CON	ICLUSIONS	7	
	REF	TERENCES		
	TAB	BLES		
	FIG	BURES		

TABLES

- I. Logs of Normalized P-P Amplitudes
- II. Intersecting Subarrays For Various Displacement Distances and Azimuths Used in Spatial Correlation
- III. Coefficients of Correlation For Individual Events At Designated Spatial Shifts
 - IV. Subarray Correlation For The Colombia Earthquakes
 - V. Correlations of Similarly Oriented Subarrays For The Colombia Earthquakes
- VI. Subarray Correlations For The Two Kazakh Explosions
- VII. Correlations of Similarly Oriented Subarrays For The Kazakh Explosions

FIGURES

- 1.
- 2. Event 152
- 3. Event 171
- 4. Event 197
- 5. Event 219
- 6. Event 238
- 7. Event 247
- 8. Event 253
- 9. Event 254
- 10. Event 291
- 11. Event 359
- 12.
- 13. N. Colombia, 21 December 1965, Seismogram No.6393,C4,Leg 1
- 14. N. Colombia, 21 December 1965, Seismogram No. 6293, C4, Leg 2
- 15. N. Colombia, 21 December 1965, Seismogram No.6393,C4 Leg 3
- 16. N. Colombia, 12 June 1966, Seismogram No. 7635, C4, Leg 1
- 17. N. Colombia, 12 June 1966, Seismogram No. 7635, C4, Leg 2
- 18. N. Colombia, 12 June 1966, Seismogram No. 7635, C4, Leg 3
- 19.
- 20.

ABSTRACT

Spatial correlations of amplitude anomalies have been conducted over LASA and LASA subarrays to test the hypothesis that these anomalies exhibit spatial stationarity. The evidence indicates that the anomaly process cannot be considered to be covariance stationary.

I. INTRODUCTION

This report describes results of part of a study of amplitude anomalies at LASA. The object of this part of the study was to see if these anomalies could be treated, in some sense, as a stochastic process.

It has been demonstrated (1,2) that the normalized short period peak-to-peak amplitudes of teleseismic events have a log normal distribution. That is, if the amplitudes of a set of events from the same geographic region are measured and their logarithms are taken, ther we find that

$$\log a_{ij} = \log L_{ij} - \frac{1}{N} \sum_{j=1}^{N} \log L_{ij},$$

has a normal distribution. In this equation, the L_{ij} are either the measured peak-to-peak amplitudes at all elements in a LASA subarray or are the peak-to-peak amplitudes observed at the center elements of the subarrays. The index j is on the seismometer and i is an event index.

The distribution of log amplitudes is not normal if the collection includes all elements in LASA. The variance of log a_{ij} is larger in the case where the L_{ij} are the observed amplitudes at the center elements. The variances are the same at each subarray when the L_{ij} are the amplitudes of the elements in a subarray. Thus, these variances can be pooled after normalization.

These anomalies are assumed to be real in that a precisely repeated event should produce the same amplitudes at the seismometers as the original. The anomalies vary however for events from the same geographic region and it is unlikely that a calibration of the earth would be a practical procedure with which to eliminate anomaly effects. Rather, a statistical approach may be a more reasonable way to proceed.

The fact that the anomalies in the subarrays can be pooled after normalization suggests that one may successfully hypothesize that these anomalies exhibit spatial stationarity. That is, although there may be slowly varying anomaly effects with distance*, with these removed the expectation of a particular amplitude anomaly is independent of spatial location.

Beyond this we desire that the anomaly process be covariance stationary. If this is so than the covariance function will serve as a measure of the distance which should be placed between seismometers so that they will exhibit independent amplitude estimates. Further, since the anomalies are log-normal no other statistic is needed since the covariance function is a complete statistic for normally distributed variables.

2. SPATIAL CORRELATIONS OVER LASA

This section investigates the possibility of correlation among the peak-to-peak amplitudes across all of LASA. Data used were from eleven Fiji Island earthquakes which occurred at 243° azimuth and from 9,500 km to 10,500 km distance from the center of LASA. From these eleven events, correlation coefficients were computed as spatial displacements were made over LASA. In the computations, the logarithms of the normalized data are used for the reasons cited in the introduction of this report. We define the estimate of the coefficient of correlation to be

$$r = (\sum X_{i}Y_{i}) / (\sum X_{i}^{2})^{\frac{1}{2}} (\sum Y_{i}^{2})^{\frac{1}{2}}, \qquad (1)$$

^{*} We note, however, that on a larger scale, these slow variations also are distributed as log normal.

i. e. we have set the mean of X and of Y to be zero. The justification for pre-determining X and Y is as follows: the estimate of the coefficient of correlation is generally defined to be

$$r = \frac{\text{Cov }(X, Y)}{\sqrt{\text{Var } X}} \sqrt{\text{Var } Y}$$

which for small samples turns out to be

$$r = \frac{(N-1) \sum (X_{1} - \bar{X}) (Y_{1} - \bar{Y})}{\sqrt{(N-1) \sum (X_{1} - \bar{X})^{2}} \sqrt{(N-1) \sum (Y_{1} - \bar{Y})^{2}}} = \frac{\sum (X_{1} - \bar{X}) (Y_{1} - \bar{Y})}{\sqrt{\sum (X_{1} - \bar{X})^{2}} \sqrt{\sum (Y_{1} - \bar{Y})^{2}}}$$
(2)

Now r is the coefficient of correlation for the two dimensional distribution of (X, Y). But the instruments that make up X and Y are actually subsets of the set, say, Z which contains the 21 subarray center seismometers. Since X and Y are samples of Z, and since we seek the mean of the population from which X and Y were drawn, it is reasonable to use Z as a better estimate of the population mean than that given by either \bar{X} or \bar{Y} . Due to the normalizing process, \bar{Z} is zero. Setting $\bar{X} = \bar{Y} = \bar{Z} = 0$ in (2) yields the equation given by (1). In addition, another degree of freedom is gained in estimating the coefficient of correlation since the mean is not estimated. We will apply equation (1) throughout the remainder of this report for all calculations of the coefficient of correlation.

A. LASA Spatial Correlations Using Average Log of Normalized Peak-To-Peak Amplitudes.

Since all eleven events were relatively closely grouped in comparison to the overall path, the average of the logarithms of the normalized peak-to-peak amplitudes at a given subarray was used as an estimate of the true value for that subarray (cf.Table I).

These average values were used in the calculations of the spatial correlation coefficients. Correlations were computed along a line parallel to the incoming signal (243° az.) and another set along a line perpendicular to the incoming signal (153° az.). Due to the configuration of LASA, few displacements exist where enough subarrays intersect to compute a valid coefficient of correlation. Table II lists the subarrays which were selected as intersecting at various displacements. Figure 1 presents the coefficients of correlation plotted against the spatial shifts using the average logarithm of the normalized peak-to-peak amplitudes over all events as the estimate of the true value for each subarray. The numerical values are shown in Table III, Part A.

B. LASA Spatial Correlations of Single Events

The correlation coefficients for the individual events were computed. That is, each event was consimpled singly as the spatial shifts were made across LASA at 153° and 243° azimuth as in A. The coefficients are presented in Table III, Part B. Figures 2 through 11 show the behavior of the coefficients as the spatial displacements are made. Note that Table III, Part C, contains the average coefficient of correlation, r, for various displacements, and Figure 12 presents the graph of r vs. the spatial shifts. It is of interest to compare the graphs of Figures 1 and 12, i.e. the coefficient of correlation of the average normalized amplitudes vs. displacement against the average coefficient of correlation of the individual events vs. displacement.

3. SPATIAL CORRELATIONS OVER LASA SUBARRAYS

After examining correlations over all LASA, we directed our attention to spatial correlations over subarrays. The concept of

spatial shifting over a subarray is the same as that over all of LASA except that unlike LASA the configuration of a subarray is well defined in terms of concentric circles and radial spokes every 60°. The spatial shifts over the subarrays were made along the legs (radial spokes) in this instance rather than with respect to the origin of the event.

A. Subarray Spatial Correlations for Single Events

Three N. Colombia events (21 Dec. 65; 21 April 66; 66) were chosen on the basis of availability of complete tapes of all 525 instruments in LASA for events which originated very close together. Again the logarithms of the normalized data were used to compute the coefficient of correlation and the means of the data samples were set to zero as motivated in Section 2, LASA. As spatial displacements were made across each subarray (at 0.5 km increments), coefficients of correlation were computed and graphs of coefficients vs. displacement were drawn for each leg for every subarray. Some representative graphs are given in Figures 13 through 18. The patterns in the graphs were cross-checked among the same subarrays for all three events and among similarly oriented subarrays for the same event. No consistent relationships were discovered with this approach. Contouring (Figures 19 and 20 are two examples) seemed to suggest certain patterns (viz., that the contour "pointed" towards the direction of the event) and some consistencies were found, but the presence of exceptional and contradictory contours made such a conclusion at best doubtful. It was thought that teleseismic explosions occurring close together might be better sources to study for contouring in this manner. This step is taken in 3-C below, but first we will mention some additional investigation of the three N. Colombia events.

B. Testing For Correlation Among Three Colombia Earthquakes
Coefficients of correlation were computed over each subarray
for paired events using the normalization described earlier and
equation (1). That is, the twenty-five readings of the A subarray
for, say, the 21 December 1965 earthquake, were tested against the
respective twenty-five readings of this subarray for, say the 21
April 1966 earthquake. The three independent earthquakes yield
three distinct pairs for testing in this manner, and each pair
of events has a maximum of twenty-one coefficients to be computed...

In nearly every case, the coefficient was significant (i.e. correlation existed) and in many cases (nearly two-thirds) the coefficients were between .8 and 1.0. Table IV summarizes the results of these calculations. This test can be considered as examining the correlation among the instrument responses as the source distance is varied. In the next step, the converse of this procedure was done ... the respective elements for similarly oriented subarrays (account being taken for long and short configurations) were inspected for correlation for each earthquake. This time, correlations were not detected with the exception of one case which may be ascribed to chance (cf. Table V).

C. Subarray Spatial Correlations For Two Shots

one coefficient corresponding to each subarray.

Two nuclear shots from Kazakh (November 21, 1965; February 13, 1966) were selected to compare with the results of the three Colombian earthquakes. The correlation coefficients were contoured as had been done of the earthquakes, and a search was made to detect similar patterns between the same subarrays for the two shots. The number of similarities found was no greater than that expected purely by chance.

D. <u>Testing For Correlation Between Two Kazakh Shots</u>

The coefficient was computed over each subarray for both events as was described in 3-B. Six of the subarrays registered a positive correlation, one a negative, and the remaining thirteen (one subarray was inoperative) no correlation, as is shown in Table VI. Performing the converse test on this data provided results similar to those obtained from the earthquakes, i.e., no correlations were detected among similarly oriented subarrays for the same event (cf Table VII).

4. CONCLUSIONS

Only tentative conclusions can be drawn from this data. The sparce sampling of the LASA array limits the reliability of the correlation coefficients which were computed. For this reason a uniformly spaced grid of seismometers would have aided this study.

It is likely that the anomaly process cannot be considered to be spatial covariance stationary. Since this process, is in fact, a description of the underlying geology one might have hypothesized this from the beginning. The author does not have a simple explanation for the log-normal distribution of the anomalies or for their apparent stationarity although this too reflects the geology.

REFERENCES

- Broome, P. W., Frankowski, D. E., and Klappenberger, F. A.,
 January, 1967, "Amplitude Anomalies at LASA", <u>Report No.</u>
 <u>LL-4</u>, Applied Research, Teledyne, Inc., Alexandria, Virginia.
- Klappenberger, F. A., 9 June, 1967, "Distribution of Short Period P-Phase Amplitudes over LASA", <u>Report No. 187</u>, Teledyne, Inc., Alexandria, Virginia.
- 3. The dates of the events are: February 4, February 8, February 17, February 26, March 10, March 12, March 20, April 16, April 25: all in 1966.

TABLE I

ogs of Normalized P-F Amplitudes

4	Number					is N						
ibarray	152	171	197	513	238	247	253	254	271	342	359	AVG.LOG.
tion Bl	086	.143	022	108	140	.057	190	- 168	033		900	, 5
B2	.065	.013	061	- 260	- 018	500	750	6.00	100	ł	900.	****
r m	190	190	000		090				-1123	ı	C7T-	440.1
3 1	100	100.	Z	1	900	151.	040	.041	041	ı	.017	.043
B 4	.100	.230	.310	027	.025	104	.295	.041	.117	ı	.204	.140
C1	013	013	097	036	.283	.146	980	134	274	-,092	ı	740
22	292	102	•		495	310	208	208	276	,	-,125	224
C3	990	.161	097	101	.107	.057	155	.065	041	ı	.228	.037
24	600	-:137	398	027	.258	.093	104	.076	.068	ı	027	.002
	4											ı
D1	.193	013	.310	.369	-408	.270	027	.204	ı	.100	ı	.202
05	244	1	301		125	.045	081	1	ı	ı	ı	141
03	.041	041	.253	600	276	1	1	ı	ı	.283	980	.051
104	229	377	•	284	018	187	086	013	032	168	071	147
		1 23			Č					9		
1 (1		1	C71.	101-	-107	ı	145	032	252	148
E2	ı	071		ı	495	252	1	585	161	0€6	125	251
E3	600	1.444		108	367		051	018	ı	420	1	200
E4	920.	.427	046	.124	.350	009	.155	.170	.230	400.	ı	.148
F1	ı	ı	ı	ť	.107	108	ı	ı	149	1	181	083
F2	.223	.230	071	102	.107	009	004	027	086	.188	980	049
E B	.352	.179	.277	.161	.204	.223	.250	.188	980.	ı	.297	.222
F4	143	215	022	600	. 1	125	161	081	.152	.025	125	070
90	990-	101	046	600	- 125	057	107	167	320	170	7.50	
							24	01.		6/11	170.	Ten.

TABLE II

Intersecting Subarrays For Various Displacement Distances and Azimuths Used In Spatial Correlation

A. Direction of Displacement - 243°

+ 10 km	+ 18 km	+ 22 km
Cl, B4	B3, D3	AO, D3
B1, A0	B1, B3	C1, C4
C2, B2	D1, B1	B1, B3
B2, C3	C1, C4	C2, C3
AO, B3	C2, C3	D1, B1
B4, C4		,

B. Direction of Displacement - 153°

<u>+ 10 km</u>	<u>+ 17 km</u>	<u>+ ?5 km</u>
B3, C4	D2, B2	D2, B4
C3, B3	B2, B1	B2, D4
C2, B1	C3, C4	D1, E1
AO, B4	B4, D4	
B1, C1	C2, C1	
<u>+</u> 80 km	+ 90 km	
E3, E4	F4, AO	
F2, D2	E1, F2	
B4, F4	C3, E2	

This table lists the intersecting subarrays for various displacement distances and azimuths which were used in the spatial correlation described in the report text.

TABLE III

Coefficients of Correlation For Individual Events At Designated Spatial Shifts

Direction of Displacement-2430

Direction of Displacement-153°

Dis	placemen	nt, km		Part A	<u>\</u>				
	<u>+</u> 10	<u>+</u> 18	<u>+</u> 22	<u>+</u> 10	<u>+</u> 17	<u>+</u> 35	<u>+</u> 80	<u>+</u> 90	
Coeff. of Correl. of Avg. Logs	.447	087	105	.173	584	687	604	128	cf.Fig 1

PART B

Thronk No.		Γ	1	T			1			
Event Nos	•		ł				ļ			
152	403	004	112		.341	401	х	950	x	
171	121	207	221		.108	829	ж	ж	826	
197	877	.057	.141		.469	.146	х	х	x	
219	389	724	753		.374	.381	x	х	x	
238	029	.224	. 395		035	530	979	x	x	cf.Figs
247	.491	.236	.236	ı	.194	689	567	x	330	2-11
253	.592	-894	.894	ľ	.227	797	273	773	x	
254	206	615	615		. 328	506	x	x	x	
271	.880	.820	.820		.085	921	x	x	.727	
359	223	686	686		066	828	x	æ	946	

PART C

Avg. Coef. of Correl. of Each Event, r		001	010	.203	498	606	861	344	cf.Fig.
---	--	-----	-----	------	-----	-----	-----	-----	---------

TABLE IV
Subarray correlations for the Colombia Earthquakes

		Dec 65 Apr 66	3	Dec 65 Jun 66		Apr 66 Jun 66
	Computed	Critical r.5%	Computed r	Critical r.5%	Computed r	Critical r,5%
B1	0.693	0.388	0.613	0.388	0.537	0.388
F3	0.955	0.388	0.851	0.388	0.776	0.388
P4	0.987	0.388	-0.477	0.388	-0.503	0.388
AO	0.410	0.388	0.637	0.388	0.235	0.388
B 3	0.815	0.388	0.754	0.388	0.959	0.388
C4	0.733	0.396	0.933	0.388	0.943	0.396
84	0.815	0.388	0.754	0.396	0.790	0.396
Ċ1	0.939	0:388	0.893	0.388	0.844	0.388
C2	0.898	0.388	0.782	0.388	0.639	0.308
B2	0.857	0.388	0.767	0.388	0.945	9.388
C3	0.902	0.388	0.746	0.388	0.836	0.388
D3	0.986	0.388	0.977	0.388	0.979	0.388
D4	0.563	0.388	0.368	0.388	0.634	0.388
Dl	0.831	0.404	0.882	0.404	0.931	0.388
D2	0.412	0.396	0.452	0.300	0.511	0.396
E3	0.945	0.388	0.815	0.388	0.843	0.388
E4	0.450	0.388	0.511	0.388	0.805	0.388
E1	0.900	0.388	0.825	0.388	0.806	0.388
F1	0.956	0.388	0.916	0.388	0.898	0.388
E2	0.873	0.388	0.889	0.388	0.901	0.388
F2	0.956	0.388	0.860	0.388	0.860	0.388

Subarray correlations for the Colombia earthquakes

The critical values are determined by using the "t" distribution where

$$t = r \left(\frac{n-1}{1-r^2} \right)^{\frac{r}{2}}$$

A detailed explanation is presented in Snedecor's "Statistical Methods" Fifth Edition, pp 173, 174.

TABLE V

Correlations of Similarly Oriented Subarrays For The Colombia Earthquakes

	21 Dec	Dec. 1965	21 Apr. 1965	. 1965	12 Jun	12 Jun. 1965
	Computed	Critical r,5%	Computed	Critical r,5%	Computed	Critical r,5%
C4, E1	0.239	0.388	0.232	0.396	0.121	0.388
В1, С2	0.082	0.388	090.0	0.388	0.257	0.388
B1, D2	0.200	0.388	-0.068	0.396	0.239	0.388
B1, E3	0.332	0.388	0.195	0.388	0.495	0.388
c2, D2	-0.137	0.388	-0.137	0.396	-0.340	0.388
C2, E3	0.005	0.388	-0.033	0.388	0.143	0.388
D2, E3	0.227	0.388	0.042	0.396	0.108	0.388

TABLE VI Subarray Correlations For The Two Kazakh Explosions

Subarray Designation	Coeff. of Correlation	Critical r, 5%
B1	0.067	0.388
F3	0.021	0.388
F4	-0.042	0.388
AO	0.615	0.388
В3	0.898	0.388
C4	0.905	0.388
C1	0.276	0.388
C2	-0.264	0.388
B2	-0.157	0.388
C3	-0.030	0.386
D3	0.275	0.388
D4	0.300	0.388
Dl	0.333	0.388
D2	0.071	0.388
E3	0.554	0.388
E4	0.788	0.388
El	0.044	0.388
Fl	-0.573	0.398
E2	0.595	0.388
F2	0.220	0.388

Subarray correlations for the two Kazakh explosions.

TABLE VII

Correlations of Similarly Oriented Subarrays
For The Kazakh Explosions

	21 Nov.	1965	13 Feb.	1966
	Computed	Critical r, 5%	Computed r	Critical r, 5%
C4, E1	-0.075	0.388	-0.062	0.388
B1, C2	0.068	0.388	0.021	0.388
B1, D2	0.159	0.388	-0.282	0.388
B1, E3	0.170	0.388	0.024	0.388
C2, D2	0.229	0.388	0.294	0.388
C2, E3	0.214	0.388	-0.175	0.388
D2, E3	-0.240	0.388	-0.117	0.388




Figure 1.




Figure 2. Event 152




Figure 3. Event 171





Figure 4. Event 197



• • •

Figure 5. Event 219




Figure 6. Event 238




Figure 7. Event 247




Figure 8. Event 253




Figure 9. Event 254




Figure 10. Event 291




Figure 11. Event 359




Figure 12.




Figure 13. N. Colombia, 21 December 1965 Seismogram No. 6393,C4,Leg 1




Figure 14. N. Colombia, 21 December 1965 Seismogram No. 6393,C4,Leg 2




Figure 15. N. Colombia, 21 December 1965 Seismogram No. 6393,C4,Leg 3




Figure 16. N. Colombia, 12 June 1966 Seismogram No. 7635, C4, Leg 1




Figure 17. N. Colombia, 12 June 1966 Seismogram No. 7635, 24, Leg 2






Figure 18. N. Colombia, 12 June 1966 Seismogram No. 7635, C4, Leg 3





Security Classification

(Security classification of title, bady of abstract or	NT CONTROL DATA - RAD
1. ORIGINATING ACTIVITY (Corporte author)	INI CONTROL DATA - RAD Ind indexing annotation must be entered when the averall report to clearified)
TELEDYNE, INC.	24 REPORT SECURITY CLASSIFICATION
ALEXANDRIA, VIRGINIA	Unclassified
I REPORT TITLE	
SPATIAL CORRELATION OF AMP	
4. DESCRIPTIVE HOTES (Type of report and Inchesive de	Diso)
Scientific	
S. AUTHOR(3) (Last name, Bret name, Initial)	
Klappenberger, F. A.	
4 REPORT DATE	76. TOTAL NO. OF PAGES 78. NO. OF HEPS 3
7 September 1967	36
SA. CONTRACT OR SRANT NO.	Se. ORIGINATOR , REPORT NUMBER(S)
F 33657-67-C-1313	
	195
VELA T/6702	
ARPA Order No. 624	Sb. OTHER REPORT HO(S) (Any other ambers that may be excipted
 ARPA Program Code 	
10. A VA IL ABILITY/I MITATION MATERIA	o special export controls and each trans
TO CONCERNING	III.S OF TOYOTAN matical
mittal to foreign governmen with prior approval of Chie	III.S OF TOYOTAN matical

Spatial correlations of amplitude anomalies have been conducted over LASA and LASA subarrays to test the hypothesis that these anomalies exhibit spatial stationarity. The evidence indicates that the anomaly process cannot be considered to be covariance stationary.

UNCLASSIFIED

14 KEY WORDS	LINK A		LINK B		LINK C	
	MOLE	WT	ROLE	WT	ROLE	WT
LASA					TE	
P-Waves			1 1			
Amplitude Anomalies	1 3		1 1			
Correlation Techniques			ļ .			
Seismometers			1 1		ļ l	
			1			
			1			
	j		1 1			
			1 1			

INSTRUCTIONS

- ORIGINATING ACTIVITY: Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.
- 2e. REPORT SECURITY CLASSIFICATION: Enter the overall security classification of the raport. Indicate whether "Restricted Dats" is included. Warking is to be in accordance with appropriete security regulations.
- 25. GROUP: Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces industrial Manual. Enter 'he group number. Also, when epplicable, show that optional markings have been used for Group 3 and Group 4 as authorized.
- 3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in ell cases should be unclessified. If a meeningful title cannot be selected without clessification, show title clessification in all cepitals in parenthesis immediately following the title.
- 4. DESCRIPTIVE NOTES: If appropriate, enter the type of report, e.g., Interim, progress, summery, annual, or final. Give the inclusive dates when a specific reporting period is covered.
- S. AUTHOR(S): Enter the name(s) of author(e) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.
- REPORT DATE: Enter the date of the report se day, month, year; or month, year. If more then one date appears on the report, use date of publication.
- 7a. TOTAL NUMBER OF PAGES: The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 76. NUMBER OF REFERENCES: Enter the total number of references cited in the report.
- Ba. CONTRACT OR GRANT NUMBER: If uppropriate, enter the applicable number of the contract or grant under which the report was written.
- 8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such se project number, subproject number, eystem numbers, tesk number, etc.
- 9a. ORIGINATOR'S REPORT NUMBER(S): Enter the officiel report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.
- 95. OTHER REPORT NUMBER(\$): If the report has been seeigned any other report numbers (either by the originator or by the aponeor), elso enter this number(s).
- 10. AVAILABILITY/LIMITATION NOTICES: Enter eny limitetione on further disseminetion of the report, other then those

impowed by accurity classification, using standard atstamente such es:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not euthorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other quelified DDC users shall request through
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users ohall request through
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through

If the report has bean furnished to the Office of Technical Services, Department of Commerca, for sain to the public, Indicate this fact and anter the prica, if known.

- IL SUPPLEMENTARY NOTES: Use for additional explana-
- 12. SPONSORING MILITARY ACTIVITY: Enter the name of the departmental project office or laboratory aponaoring (pering for) the research and development. Include address.
- 13. ABSTRACT: Enter an abstract giving a brief and fectual aummary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional apace is required, a continuation elset shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (75), (5), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. KEY WORDS: May words are technically meaningful terms or short phrases that characterize a report end may be used ee Index entries for estaloging the report. Key words must be selected so that no security classification is required. Identifiers, such as aquipment model designation, trade name, military project code name, geographic location, may be used se key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.